arXiv:0812.1637v1 [physics.gen-ph] 9 Dec 2008

ON NONLINEAR EQUATIONS CONNECTED WITH SIX-DIMENSIONAL PLEBANSKI SPACE

V.Dryuma*, L.Bogdanov**

*IMI AS RM,
5 Academiei Street, 2028 Kishinev, Moldova,

**L.D. Landau ITP RAS,
Kosygin str. 2, Moscow 119334, Russia

Abstract

An examples of Monge-Ampere equations connected with six-dimensional generalization of the Plebanski four-dimensional space are considered.

1 The four-dimensional second heavenly equation

Plebanski second heavenly equation is connected with the metric of the form

$$ds^2 = C(x,y,z,u)dz^2 + 2\,B(x,y,z,u)dz\,du + A(x,y,z,u)du^2 + dx\,du + dy\,dz \tag{1}$$

Conditions for the metric (1) to be Ricci-flat

$$R_{ik} = 0$$

lead to the equations on the coefficients

$$R_{xz} = R_{yz} = \frac{\partial}{\partial x}B(x,y,z,u) + \frac{\partial}{\partial y}C(x,y,z,u) = 0,$$

$$R_{xu} = R_{yu} = \frac{\partial}{\partial x}A(x,y,z,u) + \frac{\partial}{\partial y}B(x,y,z,u) = 0.$$

$$R_{zz} = 2\,\left(\frac{\partial}{\partial x}A(x,y,z,u)\right)\frac{\partial}{\partial x}C(x,y,z,u) + 2\,A(x,y,z,u)\frac{\partial^2}{\partial x^2}C(x,y,z,u) +$$

$$+2\,\left(\frac{\partial}{\partial x}B(x,y,z,u)\right)\frac{\partial}{\partial y}C(x,y,z,u) + 4\,B(x,y,z,u)\frac{\partial^2}{\partial x\partial y}C(x,y,z,u) + 2\,\frac{\partial^2}{\partial x\partial z}B(x,y,z,u) -$$

$$-2\,\frac{\partial^2}{\partial u\partial x}C(x,y,z,u) - 2\,\left(\frac{\partial}{\partial x}B(x,y,z,u)\right)^2 + 2\,C(x,y,z,u)\frac{\partial^2}{\partial y^2}C(x,y,z,u) -$$

$$-2\,\left(\frac{\partial}{\partial x}C(x,y,z,u)\right)\frac{\partial}{\partial y}B(x,y,z,u) = 0,$$

$$R_{zu} = 2\,A(x,y,z,u)\frac{\partial^2}{\partial x^2}B(x,y,z,u) + 2\,\left(\frac{\partial}{\partial x}B(x,y,z,u)\right)\frac{\partial}{\partial y}B(x,y,z,u) +$$

$$+4\,B(x,y,z,u)\frac{\partial^2}{\partial x\partial y}B(x,y,z,u) + \frac{\partial^2}{\partial x\partial z}A(x,y,z,u) - \frac{\partial^2}{\partial u\partial x}B(x,y,z,u) -$$

$$-2\,\left(\frac{\partial}{\partial x}C(x,y,z,u)\right)\frac{\partial}{\partial y}A(x,y,z,u) + 2\,C(x,y,z,u)\frac{\partial^2}{\partial y^2}B(x,y,z,u) -$$

$$-\frac{\partial^2}{\partial y\partial z}B(x,y,z,u) + \frac{\partial^2}{\partial u\partial y}C(x,y,z,u) = 0,$$

$$R_{uu} = 2\,A(x,y,z,u)\frac{\partial^2}{\partial x^2}A(x,y,z,u) + 4\,B(x,y,z,u)\frac{\partial^2}{\partial x\partial y}A(x,y,z,u) +$$

$$+2\,\left(\frac{\partial}{\partial y}B(x,y,z,u)\right)\frac{\partial}{\partial x}A(x,y,z,u) + 2\,\left(\frac{\partial}{\partial y}C(x,y,z,u)\right)\frac{\partial}{\partial y}A(x,y,z,u) +$$

$$+2\,C(x,y,z,u)\frac{\partial^2}{\partial y^2}A(x,y,z,u) + 2\,\frac{\partial^2}{\partial u\partial y}B(x,y,z,u) - 2\,\frac{\partial^2}{\partial y\partial z}A(x,y,z,u) -$$

$$-2\,\left(\frac{\partial}{\partial x}B(x,y,z,u)\right)\frac{\partial}{\partial y}A(x,y,z,u) - 2\,\left(\frac{\partial}{\partial y}B(x,y,z,u)\right)^2 = 0.$$

After the substitution

$$C(x,y,z,u) = -\frac{\partial^2}{\partial x^2}\theta(x,y,z,u), \quad B(x,y,z,u) = \frac{\partial^2}{\partial x\partial y}\theta(x,y,z,u),$$

$$A(x,y,z,u) = -\frac{\partial^2}{\partial y^2}\theta(x,y,z,u)$$

metric (1) takes the form

$$ds^2 = -\left(\frac{\partial^2}{\partial x^2}\theta(x,y,z,u)\right)dz^2 + 2\,\left(\frac{\partial^2}{\partial x\partial y}\theta(x,y,z,u)\right)dz\,du - \left(\frac{\partial^2}{\partial y^2}\theta(x,y,z,u)\right)du^2 + dx\,du + dy\,dz.$$

It is Ricci-flat if the function $\theta(x,y,z,u)$ satisfies the second Plebanski equation

$$\frac{\partial^2}{\partial u\partial x}\theta(x,y,z,u) + \frac{\partial^2}{\partial y\partial z}\theta(x,y,z,u) + \left(\frac{\partial^2}{\partial y^2}\theta(x,y,z,u)\right)\frac{\partial^2}{\partial x^2}\theta(x,y,z,u) - \left(\frac{\partial^2}{\partial x\partial y}\theta(x,y,z,u)\right)^2 = 0 \tag{2}$$

[1].

2 Six-dimensional generalization

We introduce the following six-dimensional generalization of the metric (1),

$$ds^2 = dx\,du + dy\,dv + dz\,dw + A(x,y,z,u,v,w)du^2 +$$

$$+2\,B(x,y,z,u,v,w)du\,dv + 2\,E(x,y,z,u,v,w)du\,dw + C(x,y,z,u,v,w)dv^2 +$$

$$+2\,H(x,y,z,u,v,w)dv\,dw + F(x,y,z,u,v,w)dw^2. \tag{3}$$

The Ricci tensor of the metric (3) has fifteen components. Nine of them are equal to zero due the conditions

$$\frac{\partial}{\partial u}E(x,y,z,u,v,w)+\frac{\partial}{\partial v}H(x,y,z,u,v,w)+\frac{\partial}{\partial w}F(x,y,z,u,v,w)=0,$$

$$\frac{\partial}{\partial u}B(x,y,z,u,v,w)+\frac{\partial}{\partial v}C(x,y,z,u,v,w)+\frac{\partial}{\partial w}H(x,y,z,u,v,w)=0,$$

$$\frac{\partial}{\partial u}A(x,y,z,u,v,w)+\frac{\partial}{\partial v}B(x,y,z,u,v,w)+\frac{\partial}{\partial w}E(x,y,z,u,v.w)=0. \tag{4}$$

This system of equation has solutions depending on arbitrary functions.

In a simplest case we have the solution

$$A(x,y,z,u,v,w)=\left(\frac{\partial^2}{\partial z^2}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial y^2}f(x,y,z,u,v,w)-\left(\frac{\partial^2}{\partial y\partial z}f(x,y,z,u,v,w)\right)^2,$$

$$C(x,y,z,u,v,w)=\left(\frac{\partial^2}{\partial x^2}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial z^2}f(x,y,z,u,v,w)-\left(\frac{\partial^2}{\partial x\partial z}f(x,y,z,u,v,w)\right)^2,$$

$$F(x,y,z,u,v,w)=\left(\frac{\partial^2}{\partial x^2}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial y^2}f(x,y,z,u,v,w)-\left(\frac{\partial^2}{\partial x\partial y}f(x,y,z,u,v,w)\right)^2,$$

$$E(x,y,z,u,v,w)=\left(\frac{\partial^2}{\partial y\partial z}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial x\partial y}f(x,y,z,u,v,w)-$$

$$-\left(\frac{\partial^2}{\partial x\partial z}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial y^2}f(x,y,z,u,v,w)$$

$$B(x,y,z,u,v,w)=\left(\frac{\partial^2}{\partial x\partial z}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial y\partial z}f(x,y,z,u,v,w)-$$

$$-\left(\frac{\partial^2}{\partial x\partial y}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial z^2}f(x,y,z,u,v,w)$$

$$H(x,y,z,u,v,w)=\left(\frac{\partial^2}{\partial x\partial z}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial x\partial y}f(x,y,z,u,v,w)-$$

$$-\left(\frac{\partial^2}{\partial y\partial z}f(x,y,z,u,v,w)\right)\frac{\partial^2}{\partial x^2}f(x,y,z,u,v,w)$$

depending on one arbitrary function.

Corresponding six-dimensional metric looks as

$${}^6ds^2=\left(\frac{\partial^2}{\partial w^2}K(\vec{x})\frac{\partial^2}{\partial v^2}K(\vec{x})-\left(\frac{\partial^2}{\partial v\partial w}K(\vec{x})\right)^2\right)dx^2+$$

$$+2\left(\frac{\partial^2}{\partial u\partial w}K(\vec{x})\frac{\partial^2}{\partial v\partial w}K(\vec{x})-\frac{\partial^2}{\partial w^2}K(\vec{x})\frac{\partial^2}{\partial u\partial v}K(\vec{x})\right)dxdy+$$

$$+\left(\frac{\partial^2}{\partial u^2}K(\vec{x})\frac{\partial^2}{\partial w^2}K(\vec{x})-\left(\frac{\partial^2}{\partial u\partial w}K(\vec{x})\right)^2\right)dy^2+$$

$$+2\left(\frac{\partial^2}{\partial v\partial w}K(\vec{x})\frac{\partial^2}{\partial u\partial v}K(\vec{x})-\frac{\partial^2}{\partial u\partial w}K(\vec{x})\frac{\partial^2}{\partial v^2}K(\vec{x})\right)dxdz+$$

$$+\left(\frac{\partial^2}{\partial v^2}K(\vec{x})\frac{\partial^2}{\partial u^2}K(\vec{x})-\left(\frac{\partial^2}{\partial u\partial v}K(\vec{x})\right)^2\right)dz^2+$$

$$+2\left(\frac{\partial^2}{\partial u\partial w}K(\vec{x})\frac{\partial^2}{\partial u\partial v}K(\vec{x})-\frac{\partial^2}{\partial u^2}K(\vec{x})\frac{\partial^2}{\partial v\partial w}K(\vec{x})\right)dzdy+$$

$$+dxdu+dydv+dzdw \tag{5}$$

where $K(\vec{x})=K(x,y,z,u,v,w)$ is an arbitrary function.

The Ricci tensor R_{ij} of the metric (5) has six components.

All equations

$$R_{ij}=0$$

after the substitution

$$K(x,y,z,u,v,w)=\phi(y+v+x,z+w+x)$$

are reduced to one equation

$$-\left(\frac{\partial^4}{\partial\xi\partial\rho^2\partial\xi}\phi(\xi,\rho)\right)\frac{\partial^2}{\partial\xi^2}\phi(\xi,\rho)+2\left(\frac{\partial^3}{\partial\xi\partial\rho\partial\xi}\phi(\xi,\rho)\right)^2-2\left(\frac{\partial^3}{\partial\rho^2\partial\xi}\phi(\xi,\rho)\right)\frac{\partial^3}{\partial\xi^3}\phi(\xi,\rho)-$$

$$-\left(\frac{\partial^2}{\partial\rho^2}\phi(\xi,\rho)\right)\frac{\partial^4}{\partial\xi^4}\phi(\xi,\rho)+2\left(\frac{\partial^2}{\partial\rho\partial\xi}\phi(\xi,\rho)\right)\frac{\partial^4}{\partial\xi^2\partial\rho\partial\xi}\phi(\xi,\rho)+$$

$$+2\left(\frac{\partial^3}{\partial\rho^2\partial\xi}\phi(\xi,\rho)\right)^2-\left(\frac{\partial^4}{\partial\rho^4}\phi(\xi,\rho)\right)\frac{\partial^2}{\partial\xi^2}\phi(\xi,\rho)-2\left(\frac{\partial^3}{\partial\rho^3}\phi(\xi,\rho)\right)\frac{\partial^3}{\partial\xi\partial\rho\partial\xi}\phi(\xi,\rho)-$$

$$-\left(\frac{\partial^2}{\partial\rho^2}\phi(\xi,\rho)\right)\frac{\partial^4}{\partial\xi\partial\rho^2\partial\xi}\phi(\xi,\rho)+2\left(\frac{\partial^2}{\partial\rho\partial\xi}\phi(\xi,\rho)\right)\frac{\partial^4}{\partial\rho^3\partial\xi}\phi(\xi,\rho)=0, \tag{6}$$

where

$$\xi=x+y+v,\quad \rho=z+x+w.$$

In compact form this equation can be rewritten as

$$\Delta\psi(\xi,\rho)=0$$

where

$$\psi=(\xi,\ \rho)=\left(\frac{\partial^2}{\partial\xi^2}\phi(\xi,\rho)\right)\frac{\partial^2}{\partial\rho^2}\phi(\xi,\rho)-\left(\frac{\partial^2}{\partial\xi\partial\rho}\phi(\xi,\rho)\right)^2$$

and

$$\Delta=\frac{\partial^2}{\partial\xi^2}+\frac{\partial^2}{\partial\rho^2}$$

is the Laplace operator.

Its solutions give Ricci-flat examples of the metric (5).

3 The Beltrami parameters

Two invariant equations defined by the first

$$\Delta\psi = g^{ij}\frac{\partial\psi}{\partial x^i}\frac{\partial\psi}{\partial x^j}$$

and the second

$$\Box\psi = g^{ij}\left(\frac{\partial^2}{\partial x^i\partial x^j} - \Gamma^k_{ij}\frac{\partial}{\partial x^k}\right)\psi$$

Beltrami parameters can be considered to investigate the properties of the metrics (5).

For the metric (5) the equation $\Box\phi = 0$ looks as

$$\begin{gathered}
\frac{\partial^2}{\partial u\partial x}\phi(\vec{x}) + \frac{\partial^2}{\partial w\partial z}\phi(\vec{x}) + \frac{\partial^2}{\partial v\partial y}\phi(\vec{x}) - \left(\frac{\partial^2}{\partial y^2}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x^2}f(\vec{x})\right)\frac{\partial^2}{\partial z^2}f(\vec{x}) + \\
+2\left(\frac{\partial^2}{\partial x\partial z}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x\partial z}f(\vec{x})\right)\frac{\partial^2}{\partial y^2}f(\vec{x}) - 2\left(\frac{\partial^2}{\partial x\partial z}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial y\partial z}f(\vec{x})\right)\frac{\partial^2}{\partial x\partial y}f(\vec{x}) + \\
+2\left(\frac{\partial^2}{\partial x\partial y}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x\partial y}f(\vec{x})\right)\frac{\partial^2}{\partial z^2}f(\vec{x}) - \left(\frac{\partial^2}{\partial x^2}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial z^2}f(\vec{x})\right)\frac{\partial^2}{\partial y^2}f(\vec{x}) + \\
+2\left(\frac{\partial^2}{\partial y\partial z}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial y\partial z}f(\vec{x})\right)\frac{\partial^2}{\partial x^2}f(\vec{x}) - 2\left(\frac{\partial^2}{\partial y\partial z}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x\partial z}f(\vec{x})\right)\frac{\partial^2}{\partial x\partial y}f(\vec{x}) - \\
-\left(\frac{\partial^2}{\partial z^2}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x^2}f(\vec{x})\right)\frac{\partial^2}{\partial y^2}f(\vec{x}) - 2\left(\frac{\partial^2}{\partial x\partial y}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x\partial z}f(\vec{x})\right)\frac{\partial^2}{\partial y\partial z}f(\vec{x}) + \\
+\left(\frac{\partial^2}{\partial z^2}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x\partial y}f(\vec{x})\right)^2 + \left(\frac{\partial^2}{\partial y^2}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial x\partial z}f(\vec{x})\right)^2 + \left(\frac{\partial^2}{\partial x^2}\phi(\vec{x})\right)\left(\frac{\partial^2}{\partial y\partial z}f(\vec{x})\right)^2 = 0. \qquad (7)
\end{gathered}$$

In a special case equation (7) after the substitution

$$\phi(\vec{x}) = f(\vec{x})$$

takes the form

$$\begin{gathered}
\frac{\partial^2}{\partial u\partial x}f(\vec{x}) + \frac{\partial^2}{\partial w\partial z}f(\vec{x}) + \frac{\partial^2}{\partial v\partial y}f(\vec{x}) - 3\left(\frac{\partial^2}{\partial y^2}f(\vec{x})\right)\left(\frac{\partial^2}{\partial x^2}f(\vec{x})\right)\frac{\partial^2}{\partial z^2}f(\vec{x}) + \\
+3\left(\frac{\partial^2}{\partial x\partial z}f(\vec{x})\right)^2\frac{\partial^2}{\partial y^2}f(\vec{x}) - 6\left(\frac{\partial^2}{\partial x\partial z}f(\vec{x})\right)\left(\frac{\partial^2}{\partial y\partial z}f(\vec{x})\right)\frac{\partial^2}{\partial x\partial y}f(\vec{x}) + \\
+3\left(\frac{\partial^2}{\partial x\partial y}f(\vec{x})\right)^2\frac{\partial^2}{\partial z^2}f(\vec{x}) + 3\left(\frac{\partial^2}{\partial y\partial z}f(\vec{x})\right)^2\frac{\partial^2}{\partial x^2}f(\vec{x}) = 0. \qquad (8)
\end{gathered}$$

After the change of variables

$$f(\vec{x}) = f(x, y, z, u, v, w) = h(x+u, v+y, w+z) = h(\eta, \xi, \rho)$$

equation (8) is reduced to the form

$$\frac{\partial^2}{\partial\eta^2}h(\eta,\xi,\rho) + \frac{\partial^2}{\partial\rho^2}h(\eta,\xi,\rho) + \frac{\partial^2}{\partial\xi^2}h(\eta,\xi,\rho) + 3\left(\frac{\partial^2}{\partial\eta\partial\rho}h(\eta,\xi,\rho)\right)^2\frac{\partial^2}{\partial\xi^2}h(\eta,\xi,\rho) -$$

$$-6\left(\frac{\partial^2}{\partial\eta\partial\rho}h(\eta,\xi,\rho)\right)\left(\frac{\partial^2}{\partial\rho\partial\xi}h(\eta,\xi,\rho)\right)\frac{\partial^2}{\partial\eta\partial\xi}h(\eta,\xi,\rho)+3\left(\frac{\partial^2}{\partial\eta\partial\xi}h(\eta,\xi,\rho)\right)^2\frac{\partial^2}{\partial\rho^2}h(\eta,\xi,\rho)-$$

$$-3\left(\frac{\partial^2}{\partial\xi^2}h(\eta,\xi,\rho)\right)\left(\frac{\partial^2}{\partial\eta^2}h(\eta,\xi,\rho)\right)\frac{\partial^2}{\partial\rho^2}h(\eta,\xi,\rho)+3\left(\frac{\partial^2}{\partial\rho\partial\xi}h(\eta,\xi,\rho)\right)^2\frac{\partial^2}{\partial\eta^2}h(\eta,\xi,\rho)=0 \quad (9)$$

or

$$\Delta h(\eta,\xi,\rho)-3\det\begin{bmatrix}\frac{\partial^2}{\partial\eta^2}h(\eta,\xi,\rho) & \frac{\partial^2}{\partial\eta\partial\xi}h(\eta,\xi,\rho) & \frac{\partial^2}{\partial\eta\partial\rho}h(\eta,\xi,\rho)\\ \frac{\partial^2}{\partial\eta\partial\xi}h(\eta,\xi,\rho) & \frac{\partial^2}{\partial\xi^2}h(\eta,\xi,\rho) & \frac{\partial^2}{\partial\rho\partial\xi}h(\eta,\xi,\rho)\\ \frac{\partial^2}{\partial\eta\partial\rho}h(\eta,\xi,\rho) & \frac{\partial^2}{\partial\rho\partial\xi}h(\eta,\xi,\rho) & \frac{\partial^2}{\partial\rho^2}h(\eta,\xi,\rho)\end{bmatrix}=0,$$

where

$$\Delta=\frac{\partial^2}{\partial\eta^2}+\frac{\partial^2}{\partial\xi^2}+\frac{\partial^2}{\partial\rho^2}$$

is a three-dimensional Laplace operator.

Solutions of equation (9) are characterized the properties of the metric (5).

4 Particular solutions of equation (6)

To obtain particular solutions of partial nonlinear differential equation

$$F(x,y,z,f_x,f_y,f_z,f_{xx},f_{xy},f_{xz},f_{yy},f_{yz},f_{xxx},f_{xyy},f_{xxy},..)=0, \quad (10)$$

a following approach can be applied.

We use parametric presentation of the functions and variables

$$f(x,y,z,s)\to u(x,t,z,s),\quad y\to v(x,t,z,s),\quad f_x\to u_x-\frac{u_t}{v_t}v_x, f_s\to\quad u_s-\frac{u_t}{v_t}v_s,$$

$$f_z\to u_z-\frac{u_t}{v_t}v_z,\quad f_y\to\frac{u_t}{v_t},\quad f_{yy}\to\frac{(\frac{u_t}{v_t})_t}{v_t},\quad f_{xy}\to\frac{(u_x-\frac{u_t}{v_t}v_x)_t}{v_t},... \quad (11)$$

where variable t is considered as parameter.

Note that conditions of the type

$$f_{xy}=f_{yx},\quad f_{xz}=f_{zx},\quad f_{xs}=f_{sx}\cdots$$

are fulfilled at the such type of presentation.

As a result instead of equation (10) one gets the relation between new variables $u(x,t,z)$ and $v(x,t,z)$ and their partial derivatives

$$\Psi(u,v,u_x,u_z,u_t,u_s,v_x,v_z,v_t,v_s...)=0. \quad (12)$$

This relation coincides with initial p.d.e. for $v(x,t,z,s)=t$ and takes more general form after presentation of the functions u,v in the form $u(x,t,z,,s)=F(\omega,\omega_t...)$, $v(x,t,z,s)=\Phi(\omega,\omega_t...)$ with some function $\omega(x,t,z,s)$.

4.1 The example. Laplace equation

Two-dimensional Laplace equation

$$\frac{\partial^2}{\partial x^2} f(x,y) + \frac{\partial^2}{\partial y^2} f(x,y) = 0 \tag{13}$$

after (u,v)-transformation with the conditions

$$u(x,t) = t\frac{\partial}{\partial t}\omega(x,t) - \omega(x,t), \quad v(x,t) = \frac{\partial}{\partial t}\omega(x,t)$$

takes the form of Monge-Ampere equation

$$-\left(\frac{\partial^2}{\partial t^2}\omega(x,t)\right)\frac{\partial^2}{\partial x^2}\omega(x,t) + \left(\frac{\partial^2}{\partial t \partial x}\omega(x,t)\right)^2 + 1 = 0. \tag{14}$$

Particular solution of this equation is

$$\omega(x,t) = A(t) + x + x^2 C(t)$$

where

$$A(t) = -1/12\, \_C1\, t^3 + \_C4$$

and

$$C(t) = -\frac{1}{\_C1\, t}.$$

The elimination of the parameter t from the relations

$$4\, y \_C1\, t^2 + \_C1^2 t^4 - 4\, x^2 = 0,$$

$$6\, f(x,y) \_C1\, t + \_C1^2 t^4 - 12\, x^2 + 6\, \_C4\, \_C1\, t + 6\, \_C1\, tx = 0$$

gives us the function

$$f(x,y) =$$

$$= 1/18\, \frac{-18\, \_C4\, \_C1 - 18\, x \_C1 + 12\sqrt{-2\, y^3 \_C1 + 2\, \_C1\, \sqrt{x^6 + 3\, x^4 y^2 + 3\, y^4 x^2 + y^6} + 6\, x^2 y \_C1}}{\_C1}$$

satisfying two-dimensional Laplace equation (13).

More general solutions of equation (13) can be also constructed from solutions of equation (14) in a similar way.

Note that such type of solutions of Laplace equation may be applied in the theory of water waves.

To construct particular solutions of equation (6)

$$-\left(\frac{\partial^4}{\partial y \partial x^2 \partial y}\phi(x,y)\right)\frac{\partial^2}{\partial x^2}\phi(x,y) + 2\left(\frac{\partial^3}{\partial x^2 \partial y}\phi(x,y)\right)^2 - 2\left(\frac{\partial^3}{\partial y \partial x \partial y}\phi(x,y)\right)\frac{\partial^3}{\partial x^3}\phi(x,y) -$$

$$-\left(\frac{\partial^2}{\partial y^2}\phi(x,y)\right)\frac{\partial^4}{\partial x^4}\phi(x,y) + 2\left(\frac{\partial^2}{\partial x \partial y}\phi(x,y)\right)\frac{\partial^4}{\partial x^3 \partial y}\phi(x,y) +$$

$$+2\left(\frac{\partial^3}{\partial y\partial x\partial y}\phi(x,y)\right)^2-\left(\frac{\partial^4}{\partial y^4}\phi(x,y)\right)\frac{\partial^2}{\partial x^2}\phi(x,y)-2\left(\frac{\partial^3}{\partial y^3}\phi(x,y)\right)\frac{\partial^3}{\partial x^2\partial y}\phi(x,y)-$$

$$-\left(\frac{\partial^2}{\partial y^2}\phi(x,y)\right)\frac{\partial^4}{\partial y\partial x^2\partial y}\phi(x,y)+2\left(\frac{\partial^2}{\partial x\partial y}\phi(x,y)\right)\frac{\partial^4}{\partial y^2\partial x\partial y}\phi(x,y)=0.$$

we use the method described above.

After the transformation of the function $\phi(x,y)$ and its derivatives in accordance with the rules (11) and substitution of corresponding expressions into equation (6) one obtains relation of the type (12).

From this relation in a simplest case

$$u(x,t)=A(t)x+x^2,\quad v(x,t)=B(t)x+x,\quad B(t)=t$$

we get the equation for the function $A(t)$

$$\left(\frac{d^4}{dt^4}A(t)\right)t^2+2\left(\frac{d^4}{dt^4}A(t)\right)t+2\frac{d^4}{dt^4}A(t)+4\left(\frac{d^3}{dt^3}A(t)\right)t+4\frac{d^3}{dt^3}A(t)+2\frac{d^2}{dt^2}A(t)=0,$$

having a general solution

$$A(t)=\_C1+\_C2\,t+\_C3\,\left(\arctan(t+1)t+\arctan(t+1)-1/2\,\ln(t^2+2\,t+2)\right)+$$

$$+\_C4\,\left(1/2\,t\ln(t^2+2\,t+2)+\ln(t^2+2\,t+2)-\arctan(t+1)t\right).\tag{15}$$

Elimination of the parameter t from the conditions

$$\phi(x,y)-(A(t)x+x^2)=0,\quad y-tx-x=0$$

with the function $A(t)$ from (15) leads to the solution of equation (6)

$$\phi(x,y)=x\_C1+\_C2\,y-\_C2\,x+\_C3\,\arctan(\frac{y}{x})y-1/2\,x\_C3\,\ln(\frac{y^2+x^2}{x^2})+$$

$$+1/2\,\_C4\,\ln(\frac{y^2+x^2}{x^2})y+1/2\,x\_C4\,\ln(\frac{y^2+x^2}{x^2})-\_C4\,\arctan(\frac{y}{x})y+\_C4\,\arctan(\frac{y}{x})x+x^2.$$

5 Particular solutions of equation (9)

Equation (9), rewritten in new notations

$$\frac{\partial^2}{\partial x^2}h(x,y,z)+\frac{\partial^2}{\partial z^2}h(x,y,z)+\frac{\partial^2}{\partial y^2}h(x,y,z)+3\left(\frac{\partial^2}{\partial x\partial z}h(x,y,z)\right)^2\frac{\partial^2}{\partial y^2}h(x,y,z)-$$

$$-6\left(\frac{\partial^2}{\partial x\partial z}h(x,y,z)\right)\left(\frac{\partial^2}{\partial y\partial z}h(x,y,z)\right)\frac{\partial^2}{\partial x\partial y}h(x,y,z)+3\left(\frac{\partial^2}{\partial x\partial y}h(x,y,z)\right)^2\frac{\partial^2}{\partial z^2}h(x,y,z)-$$

$$-3\left(\frac{\partial^2}{\partial y^2}h(x,y,z)\right)\left(\frac{\partial^2}{\partial x^2}h(x,y,z)\right)\frac{\partial^2}{\partial z^2}h(x,y,z)+3\left(\frac{\partial^2}{\partial y\partial z}h(x,y,z)\right)^2\frac{\partial^2}{\partial x^2}h(x,y,z)=0,\tag{16}$$

can be transformed into the form

$$-3\left(\frac{\partial^2}{\partial x^2}\omega(x,t,z)\right)\frac{\partial^2}{\partial z^2}\omega(x,t,z)+3\left(\frac{\partial^2}{\partial x\partial z}\omega(x,t,z)\right)^2-\left(\frac{\partial^2}{\partial t^2}\omega(x,t,z)\right)\frac{\partial^2}{\partial x^2}\omega(x,t,z)+$$

$$+1+\left(\frac{\partial^2}{\partial t\partial z}\omega(x,t,z)\right)^2+\left(\frac{\partial^2}{\partial t\partial x}\omega(x,t,z)\right)^2-\left(\frac{\partial^2}{\partial t^2}\omega(x,t,z)\right)\frac{\partial^2}{\partial z^2}\omega(x,t,z)=0, \tag{17}$$

according to the rules (11) and using the substitution

$$u(x,t,z)=t\frac{\partial}{\partial t}\omega(x,t,z)-\omega(x,t,z),\quad v(x,t,z)=\frac{\partial}{\partial t}\omega(x,t,z).$$

In particular case

$$\omega(x,t,z)=A(x^2+z^2,t)$$

equation (17) takes the form

$$-24\left(\frac{\partial^2}{\partial\xi^2}A(\xi,t)\right)\left(\frac{\partial}{\partial\xi}A(\xi,t)\right)\xi-12\left(\frac{\partial}{\partial\xi}A(\xi,t)\right)^2-4\left(\frac{\partial^2}{\partial t^2}A(\xi,t)\right)\left(\frac{\partial^2}{\partial\xi^2}A(\xi,t)\right)\xi-$$

$$-4\left(\frac{\partial^2}{\partial t^2}A(\xi,t)\right)\frac{\partial}{\partial\xi}A(\xi,t)+1+4\left(\frac{\partial^2}{\partial t\partial\xi}A(\xi,t)\right)^2\xi=0, \tag{18}$$

where $\xi=x^2+z^2$.

This type of equations meet in the theory of turbulent flow [2].

Particular solution of equation (18) is of the form

$$A(\xi,t)=B(t)+\xi\,e^{Ct},$$

where the function $B(t)$ is defined by the expression

$$B(t)=1/4\,\frac{e^{-Ct}}{C^2}-3\,\frac{e^{Ct}}{C^2}+\mathit{\_C1}\,t+\mathit{\_C2}.$$

Now elimination of the parameter t from the system of equations

$$h(x,y,z)-t\frac{\partial}{\partial t}\omega(x,t,z)+\omega(x,t,z)=0,$$

$$y-\frac{\partial}{\partial t}\omega(x,t,z)=0.$$

leads in the case $\_C1=0,\ \_C2=0,\ C=1$ to the function

$$h(x,y,z)=\left(y\sqrt{y^2-3+x^2+z^2}+y^2\right)\ln\left(\frac{y+\sqrt{y^2-3+x^2+z^2}}{-3+x^2+z^2}\right)\left(y+\sqrt{y^2-3+x^2+z^2}\right)^{-1}+$$

$$+\frac{3-x^2-y^2-z^2-\ln(2)y\sqrt{y^2-3+x^2+z^2}-\ln(2)y^2-y\sqrt{y^2-3+x^2+z^2}}{y+\sqrt{y^2-3+x^2+z^2}}$$

satisfying the equation (16).

Another type of particular solutions of equation (16) can be obtained using the reduction of equation (17) after the substitution

$$\omega(x,t,z) = A(x+t,z)$$

to the form

$$\left(\frac{\partial^2}{\partial x^2}A(x,y)\right)\frac{\partial^2}{\partial y^2}A(x,y) - \left(\frac{\partial^2}{\partial x \partial y}A(x,y)\right)^2 - 1/4 = 0, \tag{19}$$

where $A(x,y) = A(x+t,z)$.

The (u,v)-transformation of equation (19) with the condition

$$u(x,t) = t\frac{\partial}{\partial t}\omega(x,t) - \omega(x,t), \quad v(x,t) = \frac{\partial}{\partial t}\omega(x,t)$$

leads to Laplace equation for the function $\omega(x,t)$

$$4\,\frac{\partial^2}{\partial x^2}\omega(x,t) + \frac{\partial^2}{\partial t^2}\omega(x,t) = 0.$$

Its general solution has the form

$$\omega(x,t) = M(x+2\,It) + N(x-2\,It),$$

where M and N are arbitrary functions.

After the choice of the function $\omega(x,t)$ and elimination of the parameter t from equations

$$A(x,y) - (t\frac{\partial}{\partial t}\omega(x,t) - \omega(x,t) = 0, \quad y - \frac{\partial}{\partial t}\omega(x,t) = 0,$$

the function $A(x,y)$ and then the function

$$\omega(x,t,z) = A(x+t,z)$$

can be found.

Elimination of parameter t from equations

$$h(x,y,z) - t\frac{\partial}{\partial t}\omega(x,t,z) + \omega(x,t,z) = 0,$$

$$y - \frac{\partial}{\partial t}\omega(x,t,z) = 0 \tag{20}$$

with a given function $\omega(x,t,z)$ allow us to obtain particular solution of equation (16).

Let us consider an example.

We take solution of the Laplace equation of the form

$$\omega(x,t) = (x+2\,It)^2 + (x-2\,It)^3$$

or

$$\omega(x,t) = x^2 - 4\,t^2 + x^3 - 12\,xt^2 + I\left(4\,tx - 6\,tx^2 + 8\,t^3\right).$$

Its imaginary part

$$\omega_1 = 4\,tx - 6\,tx^2 + 8\,t^3$$

satisfies Laplace equation.

From the conditions

$$A(x,y) - t\frac{\partial}{\partial t}\omega_1(x,t) + \omega_1(x,t) = 0,$$

$$y - \frac{\partial}{\partial t}\omega_1(x,t) = 0$$

we get the system of equations

$$A(x,y) - 16\,t^3 = 0, \quad y - 4\,x + 6\,x^2 - 24\,t^2 = 0$$

from which we eliminate the parameter t.

As a result we find the function $A(x,y)$

$$A(x,y) = 1/18\,\sqrt{6\,y - 24\,x + 36\,x^2}y - 2/9\,\sqrt{6\,y - 24\,x + 36\,x^2}x + 1/3\,\sqrt{6\,y - 24\,x + 36\,x^2}x^2$$

satisfying equation (19).

From this function after the change of variables

$$x = x + t, \quad y = z$$

we find the function

$$\omega(x,t,z) = 1/18\,\sqrt{6\,z - 24\,x - 24\,t + 36\,x^2 + 72\,tx + 36\,t^2}\left(z - 4\,x - 4\,t + 6\,x^2 + 12\,tx + 6\,t^2\right)$$

Using this function we eliminate the parameter t from relations (20) and obtain corresponding solution

$$h(x,y,z) = -\frac{1}{54}\,\frac{\left(-2 - 3\,\sqrt{2 - 3\,z + T} + 3\,z + 9\,x\sqrt{2 - 3\,z + T}\right)T}{\sqrt{-2 + 3\,z + T}} -$$

$$-\frac{1}{54}\,\frac{-12\,z + 6\,\sqrt{2 - 3\,z + T} - 18\,x\sqrt{2 - 3\,z + T} + 4 - 9\,z\sqrt{2 - 3\,z + T} + 9\,z^2 + 27\,zx\sqrt{2 - 3\,z + T} - 36\,y^2}{\sqrt{-2 + 3\,z + T}},$$

where

$$T = \sqrt{9\,z^2 - 12\,z + 4 + 36\,y^2}.$$

6 Acknowledgements

The research was partially supported in the framework of joint Russian-Moldavian research project (Grant 08.820.08.07 RF of HCSTD ASM, Moldova, and RFBR grant 08-01-90104, Russia).

References

[1] V.S. Dryuma, On solutions of a Heavenly equations and their generalization, *ArXiv: gr-qc/0611001*, v1 31 Oct 2006, p.1-13.

[2] V. Rubtsov, Monge-Ampere geometry arising in some models of turbulent flow dynamics in two and three dimensions, *http://math.univ-angers.fr/publications/ /prepub/fichiers/00242.pdf., Novemre 2007, 1–9.*